

Gauged $\mathcal{W}$ algebras

F. Delduc¹, L. Frappat², E. Ragoucy², and P. Sorba^{1,2}

*Laboratoire de Physique Théorique ENSLAPP **

Abstract

We perform an Hamiltonian reduction on a classical $\mathcal{W}(\mathcal{G}, \mathcal{H})$ algebra, and prove that we get another $\mathcal{W}(\mathcal{G}, \mathcal{H}')$ algebra, with $\mathcal{H} \subset \mathcal{H}'$. In the case $\mathcal{G} = S\ell(n)$, the existence of a suitable gauge, called Generalized Horizontal Gauge, allows to relate in this way two $\mathcal{W}$ -algebras as soon as their corresponding $\mathcal{H}$ -algebras are related by inclusion.

ENSLAPP-AL-460/94
March 1994

*URA 14-36 du CNRS, associée à l'Ecole Normale Supérieure de Lyon, et au Laboratoire d'Annecy-le-Vieux de Physique des Particules (IN2P3-CNRS).

¹ Groupe de Lyon: ENS Lyon, 46 allée d'Italie, F-69364 Lyon Cedex 07, France.

² Groupe d'Annecy: LAPP, Chemin de Bellevue BP 110, F-74941 Annecy-le-Vieux Cedex, France.

1 Introduction

One knows today how to associate a $\mathcal{W}$ -algebra to each couple formed by a simple Lie (super)algebra $\mathcal{G}$ and one of its $Sl(2)$ embeddings [1, 2, 3, 4]. Each algebra obtained in this way appears as the symmetry algebra of a Toda theory [5], which itself can be seen, in a Lagrangian formalism, as a gauged Wess-Zumino-Witten (WZW) model [1, 6]. The supersymmetric generalization of this framework, leading to a classification of $\mathcal{W}$ -superalgebras associated to couples {simple Lie superalgebras – $OSp(1|2)$ embeddings}, is also known [7, 3]. First studied in the classical case, the method has also been extended to the quantum case [8, 9, 10].

Considering the large class of $\mathcal{W}$ -algebras and superalgebras so obtained, there is no doubt that several properties of simple Lie (super)algebras could be used to relate among them different $\mathcal{W}$ -(super)algebras. Simplifying their classification, this will shed some light on their structure and motivate applications. For example, the technics of folding, allowing to relate, via symmetry of their Dynkin diagrams, simply laced algebras with non simply laced ones, has been adapted to the case of $\mathcal{W}$ -(super)algebras [11]. Therefore, it is possible, at least at the classical level, to deduce the $\mathcal{W}$ families associated to the orthogonal or symplectic (orthosymplectic in the supersymmetric case) series from those relative to the unitary ones.

In this paper, connections among $\mathcal{W}$ -algebras relative to the same $\mathcal{G}$ but different $Sl(2)$ embeddings will also be established. At this point, let us recall that any $Sl(2)$ embedding of a simple Lie algebra $\mathcal{G}$ can be seen, up to some exceptions in the D_n and E_n cases, as the principal $Sl(2)$ of at least one regular subalgebra $\mathcal{H}$ of $\mathcal{G}$. We use $\mathcal{W}(\mathcal{G}, \mathcal{H})$ to denote the corresponding $\mathcal{W}$ -algebra. As we will see in the following, at least for $\mathcal{G} = Sl(n)$, the algebra $\mathcal{W}(\mathcal{G}, \mathcal{H}') = \mathcal{W}'$ can be constructed from $\mathcal{W}(\mathcal{G}, \mathcal{H}) = \mathcal{W}$, with $\mathcal{H} \subset \mathcal{H}'$, by imposing a set of constraints $\mathcal{C}(\mathcal{W} \rightarrow \mathcal{W}')$ on the W generators and performing gauge transformations associated to these constraints. Thus we perform an Hamiltonian reduction in the same way that the $\mathcal{W}$ -algebra itself is obtained by constraining some $\mathcal{G}$ Kac-Moody currents and determining quantities which are invariant under the gauge group (G -subgroup) relative to the corresponding set of constraints $\mathcal{C}(\mathcal{G} \rightarrow \mathcal{W})$. It can appear natural to call *secondary reduction* such a procedure which allows to build the algebra $\mathcal{W}'$ from the algebra $\mathcal{W}$ by gauging a part of this last one.

We start this letter by a rapid summary on Dirac constraints theory applied to the determination of $\mathcal{W}$ -algebras (section 2). Then, we show in section 3 that the reduction $\mathcal{G} \rightarrow \mathcal{W}(\mathcal{G}, \mathcal{H}') \equiv \mathcal{W}'$ can be decomposed into two steps: $\mathcal{G} \rightarrow \mathcal{W}(\mathcal{G}, \mathcal{H}) \equiv \mathcal{W}$ followed by $\mathcal{W}(\mathcal{G}, \mathcal{H}) \rightarrow \mathcal{W}(\mathcal{G}, \mathcal{H}')$ with $\mathcal{H} \subset \mathcal{H}'$, as soon as the set of all constraints $\mathcal{C}(\mathcal{G} \rightarrow \mathcal{W})$ is included into the set $\mathcal{C}(\mathcal{G} \rightarrow \mathcal{W}')$. For $\mathcal{G} = Sl(n)$, we introduce the Generalized Horizontal Gauge (section 4) which insures that $\mathcal{W}(\mathcal{G}, \mathcal{H})$ can be related to $\mathcal{W}(\mathcal{G}, \mathcal{H}')$ as soon as $\mathcal{H} \subset \mathcal{H}'$. We illustrate our results in the $Sl(4)$ case (section 5) before concluding.

2 Constraints and W algebras: a reminder

The construction of $\mathcal{W}$ -algebras via Hamiltonian reduction has been extensively discussed in several papers [1, 6]; therefore, we will limit ourselves introducing the main tools necessary for our purpose.

We start with the Lie algebra $\mathcal{G}$ of the non compact group G . Let t^a , $a = 1, \dots, \dim \mathcal{G}$, be a matrix realization of a $\mathcal{G}$ -basis; we denote by $J = J_a t^a$ the matrix current we want to constrain. The J_a can be seen either as the right $J_+(x_+)$ or as the left $J_-(x_-)$ conserved currents, i.e. $\partial_\pm(J_\mp) = 0$ of the WZW action $S(g)$ with $g \in G$.

Let $M_0, M_\pm$ be the generators of the $S\ell(2)$ subalgebra of $\mathcal{G}$ chosen for the reduction. There exists at least one regular $\mathcal{G}$ subalgebra (we discard the exceptions arising in the D_n and E_n cases) $\mathcal{H}$ for which this $S\ell(2)$ is principal. The gradation on $\mathcal{G}$ induced by M_0 leads to the decomposition:

$$\mathcal{G} = \oplus_k \mathcal{G}_k = \mathcal{G}_- \oplus \mathcal{G}_0 \oplus \mathcal{G}_+ \quad (2.1)$$

with obviously $M_\pm \in \mathcal{G}_{\pm 1}$.

Now we demand the $\mathcal{G}_-$ part of J to satisfy the following conditions:

$$J|_{\mathcal{G}_-} = M_- \quad (2.2)$$

which in particular implies for the J -components associated to elements of $\mathcal{G}_{<-1}$, if they exist, to be zero.

The conditions (2.2) are first class constraints² $C_A^{(1)} = 0$ with $A = 1, \dots, \dim \mathcal{G}_-$, and generate through the Poisson Brackets (PB) a gauge group with Lie algebra $\mathcal{G}_+$.

The PB algebra of the current components is given by:

$$\{J_a(z), J_b(w)\}_{PB} = f_{ab}^c J_c(w) \delta(z - w) + k \delta_{ab} \delta'(z - w) \quad (2.3)$$

where f_{ab}^c are the structure constants for a given basis of $\mathcal{G}$. Moreover, the gauge transformations associated to the first class constraints $C_A^{(1)}$ act on J_a as

$$\begin{aligned} J_a(w) \rightarrow J_a(w) + \int dz \epsilon^A(z) \{C_A^{(1)}(z), J_a(w)\}_{PB} + \\ + \frac{1}{2!} \int dy \int dz \epsilon^A(z) \epsilon^B(y) \{C_B^{(1)}(y), \{C_A^{(1)}(z), J_a(w)\}\} + \dots \end{aligned} \quad (2.4)$$

where $\epsilon^A(x)$ is the parameter associated to $C_A^{(1)}(x)$. As usual in Dirac constraints theory [12], the condition $C_A^{(1)}(x) = 0$ has to be imposed after working out a PB , that is why the name *weak equation* is often used.

The bracket of two W elements can be computed *à la Dirac*. From this point of view, we have to take into account the second class constraints determined by the gauge fixing. Let us denote $\{C_a(z)\}$ the set of all (first and second class) constraints: its dimension is even and equal to $2p$, where p is also the dimension of the subgroup of $\mathcal{G}$ which has been gauged. We also introduce:

$$C_{AB}(x, y) = \{C_A(x), C_B(y)\}_{PB} \quad (2.5)$$

²In the case of half-integral grading, some of the constraints we impose may be second class, but this will not change the reasoning and conclusion hereafter.

and its inverse $C^{AB}(x, y)$ such that:

$$\int dy C_{AB}(x, y) C^{BD}(y, t) = \delta_A^D \delta(x - t) \quad (2.6)$$

The (Dirac) brackets for the $\mathcal{W}(\mathcal{G}, \mathcal{H})$ algebra read:

$$\{J_i(z), J_j(w)\}_D = \{J_i(z), J_j(w)\}_{PB} - \int dx dy \{J_i(z), C_A(x)\}_{PB} C^{AB}(x, y) \{C_B(y), J_j(w)\}_{PB} \quad (2.7)$$

In the following we will drop the z dependence, and forget the index PB (while keeping the D one). Thus, we will write

$$C_{AB} C^{BD} = \delta_A^D \quad (2.8)$$

$$\{J_i, J_j\}_D = \{J_i, J_j\} - \{J_i, C_A\} C^{AB} \{C_B, J_j\} \quad (2.9)$$

instead of (2.6) and (2.7).

3 Secondary reduction

The following property will be specially usefull for our purpose:

Property 1 *Let $\mathcal{W}(\mathcal{G}, \mathcal{H}')$ obtained from $\mathcal{G}$ by imposing the set $\{C_A\}$ of first and second class constraints. Let $\mathcal{W}(\mathcal{G}, \mathcal{H})$ a second $\mathcal{W}$ -algebra with $\mathcal{H} \subset \mathcal{H}'$ and such that the corresponding set of constraints $\{C_\alpha\}$ is a subset of $\{C_A\}$. Then, by imposing on $\mathcal{W}(\mathcal{G}, \mathcal{H})$ the constraints $\{C_{\bar{\alpha}}\}$ such that $\{C_A\} = \{C_\alpha\} \cup \{C_{\bar{\alpha}}\}$, we recover the algebra $\mathcal{W}(\mathcal{G}, \mathcal{H}')$.*

This property can be sumarized in the diagram:

$$\begin{array}{ccccccc} \mathcal{G} & \rightarrow & & \rightarrow & \{C_A\} & \rightarrow & \rightarrow & \rightarrow & \mathcal{W}(\mathcal{G}, \mathcal{H}') \\ & \searrow & & & & & & \nearrow & \\ & & \{C_\alpha\} & & & & \{C_{\bar{\alpha}}\} & & \\ & & & \searrow & & \nearrow & & & \\ & & & & \mathcal{W}(\mathcal{G}, \mathcal{H}) & & & & \end{array} \quad (3.1)$$

$$\text{with } \{C_A\} = \{C_\alpha\} \cup \{C_{\bar{\alpha}}\}$$

This property is a general feature of the Dirac Brackets (see e.g. [13]). We will show hereafter that it can be applied to the case of $\mathcal{W}$ -algebras.

Let us remark that the number of free fields in $\mathcal{W}(\mathcal{G}, \mathcal{H}')$ and in the algebra obtained by reducing $\mathcal{W}(\mathcal{G}, \mathcal{H})$ via $\{C_{\bar{\alpha}}\}$ -constraints is the same, since the number of constraints imposed in both paths is identical. Let us denote by $\{\cdot, \cdot\}_D$ the Dirac bracket associated to the reduction $\mathcal{G} \rightarrow \mathcal{W}(\mathcal{G}, \mathcal{H}')$; $\{\cdot, \cdot\}_{D1}$ those associated to $\mathcal{G} \rightarrow \mathcal{W}(\mathcal{G}, \mathcal{H})$ and $\{\cdot, \cdot\}_{D2}$ those constructed from $\mathcal{W}(\mathcal{G}, \mathcal{H})$ by imposing the constraints $\{C_{\bar{\alpha}}\}$.

To carry on the proof, let us add the following notations: in $\mathcal{W}(\mathcal{G}, \mathcal{H})$ the free fields will be $\{W_i, W_{\bar{\alpha}}\} \equiv \{W_a\}$, and Dirac brackets will read:

$$\{W_a, W_b\}_{D1} = \{W_a, W_b\} - \{W_a, C_\alpha\} \Gamma^{\alpha\beta} \{C_\beta, W_b\} \quad (3.2)$$

where we have defined

$$\Gamma_{\alpha\beta} = \{C_\alpha, C_\beta\} = C_{\alpha\beta} \quad (3.3)$$

and

$$\Gamma^{\alpha\beta} \quad \text{such that} \quad \Gamma_{\alpha\beta} \Gamma^{\beta\gamma} = \delta_\alpha^\gamma \quad (3.4)$$

Now we want to constrain the $\mathcal{W}(\mathcal{G}, \mathcal{H})$ elements which correspond to the fields $W_{\bar{\alpha}}$. This means that we introduce new brackets on $\mathcal{W}(\mathcal{G}, \mathcal{H})$:

$$\{W_i, W_j\}_{D2} = \{W_i, W_j\}_{D1} - \{W_i, C_{\bar{\alpha}}\}_{D1} \Delta^{\bar{\alpha}\bar{\beta}} \{C_{\bar{\beta}}, W_j\}_{D1} \quad (3.5)$$

with

$$\Delta_{\bar{\alpha}\bar{\beta}} = \{W_{\bar{\alpha}}, W_{\bar{\beta}}\}_{D1} \quad ; \quad W_{\bar{\alpha}} \equiv C_{\bar{\alpha}} \quad \text{and} \quad \Delta_{\bar{\alpha}\bar{\beta}} \Delta^{\bar{\beta}\bar{\gamma}} = \delta_{\bar{\alpha}}^{\bar{\gamma}} \quad (3.6)$$

The property 1 will be proved if one shows that $\{W_i, W_j\}_D = \{W_i, W_j\}_{D2}$ for any couple of elements (W_i, W_j) in $\mathcal{W}(\mathcal{G}, \mathcal{H})$ –or in the $\{C_{\bar{\alpha}}\}$ -constrained $\mathcal{W}(\mathcal{G}, \mathcal{H})$ algebra.

Let us develop (3.5) using (3.5)

$$\begin{aligned} \{W_i, W_j\}_{D2} &= \{W_i, W_j\} - \{W_i, C_\alpha\} \Gamma^{\alpha\beta} \{C_\beta, W_j\} + \\ &\quad - \left(\{W_i, C_{\bar{\alpha}}\} - \{W_i, C_\alpha\} \Gamma^{\alpha\beta} \{C_\beta, C_{\bar{\alpha}}\} \right) \Delta^{\bar{\alpha}\bar{\beta}} \left(\{C_{\bar{\beta}}, W_j\} - \{C_{\bar{\beta}}, C_\gamma\} \Gamma^{\gamma\sigma} \{C_\sigma, W_j\} \right) \\ &= \{W_i, W_j\} - \{W_i, C_\alpha\} \left(\Gamma^{\alpha\beta} - \Gamma^{\alpha\gamma} C_{\gamma\bar{\alpha}} \Delta^{\bar{\alpha}\bar{\beta}} C_{\bar{\beta}\sigma} \Gamma^{\sigma\beta} \right) \{C_\beta, W_j\} + \\ &\quad + \{W_i, C_{\bar{\alpha}}\} \Delta^{\bar{\alpha}\bar{\beta}} C_{\bar{\beta}\beta} C^{\beta\alpha} \{C_\alpha, W_j\} + \{W_i, C_\alpha\} \Gamma^{\alpha\beta} C_{\beta\bar{\beta}} \Delta^{\bar{\beta}\bar{\alpha}} \{C_{\bar{\beta}}, W_j\} + \\ &\quad - \{W_i, C_{\bar{\alpha}}\} \Delta^{\bar{\alpha}\bar{\beta}} \{C_{\bar{\beta}}, W_j\} \end{aligned}$$

This last expression can, hopefully, be simplified owing to the following identities:

$$C^{\alpha\beta} = \Gamma^{\alpha\beta} - \Gamma^{\alpha\gamma} C_{\gamma\bar{\gamma}} C^{\bar{\gamma}\beta} \quad (3.7)$$

$$C^{\alpha\bar{\alpha}} = -\Gamma^{\alpha\gamma} C_{\gamma\bar{\gamma}} C^{\bar{\gamma}\bar{\alpha}} \quad (3.8)$$

$$C^{\bar{\alpha}\alpha} = -C^{\bar{\alpha}\bar{\gamma}} C_{\bar{\gamma}\gamma} \Gamma^{\gamma\alpha} \quad (3.9)$$

which are obtained by multiplying by $\Gamma^{\alpha\beta}$ both sides (2.8) with the couple (A,D) being formed with one or two non bared greek indices. We will also need the relations

$$\Delta_{\bar{\alpha}\bar{\beta}} = C_{\bar{\alpha}\bar{\beta}} - C_{\bar{\alpha}\alpha} \Gamma^{\alpha\beta} C_{\beta\bar{\beta}} \quad (3.10)$$

and

$$\Delta^{\bar{\alpha}\bar{\beta}} = C^{\bar{\alpha}\bar{\beta}} \quad (3.11)$$

Then,

$$\begin{aligned} \{W_i, W_j\}_{D2} &= \{W_i, W_j\} - \{W_i, C_\alpha\} C^{\alpha\beta} \{C_\beta, W_j\} - \{W_i, C_{\bar{\alpha}}\} C^{\bar{\alpha}\bar{\beta}} \{C_{\bar{\beta}}, W_j\} + \\ &\quad - \{W_i, C_\alpha\} C^{\alpha\bar{\beta}} \{C_{\bar{\beta}}, W_j\} - \{W_i, C_{\bar{\alpha}}\} C^{\bar{\alpha}\beta} \{C_\beta, W_j\} \\ &= \{W_i, W_j\} - \{W_i, C_A\} C^{AB} \{C_B, W_j\} \\ &= \{W_i, W_j\}_D \end{aligned}$$

The simplest example of such a secondary reduction is the gauging of the algebra $W_3^{(2)}$ also called Bershadsky-Polyakov algebra in order to obtain the W_3 or Zamolodchikov algebra. This case has been studied in [14]. We consider other examples based on $S\ell(4)$ in section 5. Before that, let us discuss the general case of $\mathcal{W}[S\ell(n-1), \mathcal{H}]$ algebras.

4 A generalized horizontal gauge for $\mathcal{W}[Sl(n-1), \mathcal{H}]$ algebras

As developed in section 3, a sufficient condition for obtaining the $\mathcal{W}(\mathcal{G}, \mathcal{H}')$ algebra from the $\mathcal{W}(\mathcal{G}, \mathcal{H})$ one is the inclusion of the constraints relative to $\mathcal{W}(\mathcal{G}, \mathcal{H})$ in those relative to $\mathcal{W}(\mathcal{G}, \mathcal{H}')$. For $\mathcal{G} = Sl(n)$, we introduce a general gauge which guarantees the gauge passing $\mathcal{W}(\mathcal{G}, \mathcal{H}) \rightarrow \mathcal{W}(\mathcal{G}, \mathcal{H}')$ as soon as $\mathcal{H} \subset \mathcal{H}'$.

One knows that in $Sl(n)$, any $Sl(2)$ -subalgebra is principal in a regular subalgebra which itself is of the form

$$\mathcal{H} = \oplus_{i=1}^k Sl(n_i) \quad \text{with} \quad \left(\sum_{i=1}^k n_i \right) \leq n \quad (4.1)$$

Note that one may have $n_i = n_j$ while $i \neq j$.

Considering a $n \times n$ matrix representing the general element of the algebra, let us start by fixing a block diagonal part made with each $Sl(n_i)$ algebra of $\mathcal{H}$. We complete it by an $Gl(p)$ subalgebra, where $p = n - \sum_{i=1}^k n_i$, and by $(k-1)$ $U(1)$ generators, commuting with each $Sl(n_i)$ and $Gl(p)$ part. On each $Sl(n_i)$ block, there will be 1 on each entry of the minor diagonal below the main one and 0 everywhere else, except on the first row where will stand n_i different W fields. In the last $p \times p$ matrix will stand $(p^2 - 1)$ different W fields, as well as on the diagonal p -times the same W -field. This last field depends on the others k fields on the main diagonal, since the trace of an $Sl(n)$ matrix is zero. Note that the fields relative to the commutant of $\mathcal{H}$, constituted by $Gl(p)$ and the $(k-1)$ $U(1)$, generate the Kac-Moody part (spin-one fields) of the $\mathcal{W}(\mathcal{G}, \mathcal{H})$ algebra.

Now, let us decompose the rest of the $Sl(n)$ matrix into $n_i \times n_j$ ($i \neq j$), $n_i \times p$ and $p \times n_i$ submatrices. In each matrix of this type, above the block diagonal part, the W fields will show up only on the first row, the rest of the matrix entries being 0. There will be p different fields for a matrix $n_i \times p$ or $p \times n_i$. For a $n_i \times n_j$ matrix, the number of different fields will be $\min(n_i, n_j)$; when $n_j \leq n_i$ the first row will then be filled by these n_j fields, but when $n_j > n_i$, the first row will be made with $n_j - n_i$ zeros followed by the W -fields in position $n_j - n_i + 1$, $n_j - n_i + 2$, $\dots$, n_j . Below the diagonal part, one uses the same submatrix decomposition. The number of different W fields in each case will be the same as above, except that now they will occupy the last column with again p fields for the matrix $p \times n_i$ or $n_i \times p$ and for the $n_i \times n_j$ matrix, $\min(n_j, n_i)$ fields put in the first, second, up to $\min(n_j, n_i)$ -th row.

In the case $\mathcal{H}=\mathcal{G}$, the $(n-1)$ different W fields of the Abelian $\mathcal{W}(\mathcal{G}, \mathcal{G})$ algebra will occupy the second, third, up to n -th entries of the first row³, the rest of the matrix being made of 0 and 1: then we recognize the so-called horizontal gauge. By extension, we will call the gauge described above the Generalized Horizontal Gauge (G.H.G.).

We also define the Ordered Generalized Horizontal Gauge (O.G.H.G.) by disposing in the $Sl(n)$ matrix the different $Sl(n_i)$ of $\mathcal{H}$ in decreasing order, i.e. $Sl(n_i)$ will show up before

³In this special case, there is *a priori* one field on the diagonal (on the first row) but it is set to zero by the traceless condition.

$S\ell(n_j)$ if $n_i > n_j$. This gauge is such that the first row (or column) of each submatrix is completely filled with W fields (no zero: see for instance the matrix 4.2).

Let us illustrate this construction in a particular case: $\mathcal{G} = S\ell(11)$ and $\mathcal{H} = S\ell(3) \oplus S\ell(2) \oplus S\ell(2)$. Then $S\ell(p) = S\ell(4)$. One can compute that the corresponding $\mathcal{W}$ -algebra will be generated by 58 fields, the Kac-Moody part being made by $S\ell(4)$ and three $U(1)$ (We recall that the total $S\ell(11)$ matrix is traceless). In the matrix just below, we have explicitly written the entries that correspond to a 1 or a W field: the rest of the matrix is filled by zeros. The positions of the W -fields is specified by a star (*).

$$\left(\begin{array}{ccc|cc|cc|cccc} * & * & * & * & * & * & * & * & * & * \\ 1 & & & & & & & & & \\ & 1 & & & & & & & & \\ \hline & & * & * & * & * & * & * & * & * \\ & & * & 1 & & & & & & \\ \hline & & * & & * & * & * & * & * & * \\ & & * & & * & 1 & & & & \\ \hline & & * & & * & & * & * & * & * \\ & & * & & * & & * & * & * & * \\ & & * & & * & & * & * & * & * \\ & & * & & * & & * & * & * & * \end{array} \right) \quad (4.2)$$

In the matrix (4.2), the horizontal and vertical lines are drawn to make apparent the submatrices $n_i \times p$, $p \times n_i$ and $n_i \times n_j$ described above. Note that this matrix corresponds to an O.G.H.G.: as a consequence, the lines of "stars" are "full".

Now it is rather obvious that the constraints associated to the O.G.H.G. of the couple $(\mathcal{G}, \mathcal{H})$ will be included into the constraints associated to the G.H.G. of $(\mathcal{G}, \mathcal{H}')$ when $\mathcal{H} \subset \mathcal{H}'$. Thus, with the help of property 1, we conclude that we can gauge $\mathcal{W}(\mathcal{G}, \mathcal{H})$ to get $\mathcal{W}(\mathcal{G}, \mathcal{H}')$. Note that although we have started from an ordered G.H.G, after reduction, the new gauge may be not ordered anymore.

Let us take the opportunity of matrix (4.2) to treat explicitly an example. We choose, for example, to reduce the given algebra $\mathcal{W}(\mathcal{G}, \mathcal{H})$ with $\mathcal{G} = S\ell(11)$ and $\mathcal{H} = S\ell(3) \oplus S\ell(2) \oplus S\ell(2)$ up to $\mathcal{W}(\mathcal{G}, \mathcal{H}')$ with $\mathcal{H}' = S\ell(3) \oplus S\ell(4)$. Then the W -fields in position (1,4), (6,5), (6,6), ..., (6,11) will disappear as well as those in position (7,3), (7,5) and also those in position (8,5), (9,5), (10,5), (11,5). With the same conventions as in (4.2), the gauged matrix becomes:

$$\left(\begin{array}{ccc|ccc|cccc}
* & * & * & & * & * & * & * & * & * & * \\
1 & & & & & & & & & & \\
& 1 & & & & & & & & & \\
\hline
& & * & * & * & * & * & * & * & * & \\
& & * & 1 & & & & & & & \\
& & * & & 1 & & & & & & \\
& & & & & 1 & & & & & \\
\hline
& & * & & & * & * & * & * & * & \\
& & * & & & & * & * & * & * & \\
& & * & & & & * & * & * & * & \\
& & * & & & & * & * & * & * &
\end{array} \right) \quad (4.3)$$

As announced, the new gauge is a non-ordered G.H.G., and some lines of "stars" are "half-empty". The inclusion of the constraints appears naturally, and one can check independently that we have the right number of $\mathcal{W}(\mathcal{G}, \mathcal{H}')$ fields, i.e. 44.

5 An example: reduction of $\mathcal{W}(Sl(4), \mathcal{H})$ -algebras.

As an illustrative example of the developed method, let us consider the Lie algebra $Sl(4)$, which leads to four different $\mathcal{W}$ -algebras by Hamiltonian reduction, namely $\mathcal{W}[Sl(4), Sl(2)]$, $\mathcal{W}[Sl(4), 2Sl(2)]$, $\mathcal{W}[Sl(4), Sl(3)]$ (non Abelian cases) and $\mathcal{W}[Sl(4)]$ (Abelian case). Among them, the model corresponding to $\mathcal{W}[Sl(4), Sl(2)]$ is the less constrained one. Thus, let us start with this algebra. Using classical arguments, we directly deduce that a basis of W fields is given by one spin 2, four spin 1 and four spin 3/2 primary fields. Following section 4, one chooses the generalized horizontal gauge:

$$J = \left(\begin{array}{cc|cc}
2W_1 & W_2 & W_{3/2}^{+-} & W_{3/2}^{++} \\
1 & 0 & 0 & 0 \\
\hline
0 & W_{3/2}^{-+} & -W_1 + W_1^0 & W_1^+ \\
0 & W_{3/2}^{--} & W_1^- & -W_1 - W_1^0
\end{array} \right) \quad (5.1)$$

Notice that in this gauge the fields are not primary. The KM part of the $\mathcal{W}[Sl(4), Sl(2)]$ algebra is generated by the four spin one fields which form the algebra $Sl(2)^{(1)} \oplus U(1)^{(1)}$.

From the property 1 and the choice of the gauge, we deduce that one can go from the $\mathcal{W}[Sl(4), Sl(2)]$ algebra to the three others by a suitable set of constraints. More precisely, setting as (first class) constraint the condition $W_1^- = 1$, the associated gauge transformation

$$W(x) \longrightarrow \widehat{W}(x) = W(x) + \int dz \{ \alpha(z) W_1^-(z), W(x) \} + \dots \quad (5.2)$$

leads to the following transformed fields:

$$\begin{aligned}
\widehat{W}_1 &= W_1 & \widehat{W}_1^0 &= W_1^0 + \alpha \\
\widehat{W}_1^+ &= W_1^+ - 2\alpha W_1^0 + \partial\alpha - \alpha^2 & \widehat{W}_2 &= W_2 \\
\widehat{W}_{3/2}^{+-} &= W_{3/2}^{+-} & \widehat{W}_{3/2}^{++} &= W_{3/2}^{++} - \alpha W_{3/2}^{+-} \\
\widehat{W}_{3/2}^{--} &= W_{3/2}^{--} & \widehat{W}_{3/2}^{-+} &= W_{3/2}^{-+} + \alpha W_{3/2}^{--}
\end{aligned} \tag{5.3}$$

The gauge fixing is determined by imposing $\widehat{W}_1^0 = -W_1$ i.e. $\alpha = -W_1^0 - W_1$. The gauge fixed fields will read

$$\begin{aligned}
\widehat{W}_1 &= W_1 & \widehat{W}_1^+ &= W_1^+ + W_1^0 \cdot W_1^0 - W_1 \cdot W_1 - \partial W_1^0 - \partial W_1 \\
\widehat{W}_{3/2}^{+-} &= W_{3/2}^{+-} & \widehat{W}_{3/2}^{++} &= W_{3/2}^{++} + W_1^0 \cdot W_{3/2}^{+-} + W_1 \cdot W_{3/2}^{+-} \\
\widehat{W}_{3/2}^{--} &= W_{3/2}^{--} & \widehat{W}_{3/2}^{-+} &= W_{3/2}^{-+} - W_1^0 W_{3/2}^{--} - W_1 \cdot W_{3/2}^{--} \\
\widehat{W}_2 &= W_2
\end{aligned} \tag{5.4}$$

The corresponding current matrix $\widehat{J}$ is then

$$\widehat{J} = \left(\begin{array}{cc|cc} 2\widehat{W}_1 & \widehat{W}_2 & \widehat{W}_{3/2}^{+-} & \widehat{W}_{3/2}^{++} \\ 1 & 0 & 0 & 0 \\ \hline 0 & \widehat{W}_{3/2}^{-+} & -2\widehat{W}_1 & \widehat{W}_1^+ \\ 0 & \widehat{W}_{3/2}^{--} & 1 & 0 \end{array} \right) \tag{5.5}$$

The commutation relations among the $\widehat{W}$ fields can be computed explicitly using the relations (5.4) and the commutation relations of $\mathcal{W}[S\ell(4), S\ell(2)]$. One finds (and this can be checked also with the form of the $\widehat{J}$ current) that the $\widehat{W}$ fields generate the algebra $\mathcal{W}[S\ell(4), 2S\ell(2)]$, with four spin 2 and three spin 1 fields. The primary fields are given by

$$\begin{aligned}
\text{stress-energy tensor} & \quad \widehat{W}_2 + \widehat{W}_1^+ + 4\widehat{W}_1 \cdot \widehat{W}_1 \\
\text{spin 1 fields} & \quad \widehat{W}_1, \quad \widehat{W}_{3/2}^{+-}, \quad \widehat{W}_{3/2}^{--} \\
\text{spin 2 fields} & \quad \widehat{W}_1^+ - \widehat{W}_2 + 2\partial\widehat{W}_1, \quad \widehat{W}_{3/2}^{++} - \frac{1}{2}\partial\widehat{W}_{3/2}^{+-}, \quad \widehat{W}_{3/2}^{-+} + \frac{1}{2}\partial\widehat{W}_{3/2}^{--}
\end{aligned} \tag{5.6}$$

Now, one can perform a secondary reduction on the algebra $\mathcal{W}[S\ell(4), 2S\ell(2)]$ with the choice of (first class) constraints $\widehat{W}_{3/2}^{-+} = 1$ and $\widehat{W}_{3/2}^{--} = 0$, with the associated gauge transformation

$$\widehat{W}(x) \longrightarrow \widetilde{W}(x) = \widehat{W}(x) + \int dz \{ (\alpha \widehat{W}_{3/2}^{-+} + \beta \widehat{W}_{3/2}^{--})(z), \widehat{W}(x) \} + \dots \tag{5.7}$$

The transformed fields are:

$$\begin{aligned}
\widetilde{W}_1 &= \widehat{W}_1 + \alpha/2 \\
\widetilde{W}_1^+ &= \widehat{W}_1^+ - \beta \\
\widetilde{W}_2 &= \widehat{W}_2 - 4\alpha\widehat{W}_1 + 2\partial\alpha + \beta - \alpha^2
\end{aligned} \tag{5.8}$$

and more complicated expressions for $\widetilde{W}_{3/2}^{+-}$ and $\widetilde{W}_{3/2}^{++}$. The gauge fixing is obtained by imposing $\widetilde{W}_1 = 0$ and $\widetilde{W}_1^+ = 0$ that is $\alpha = -2\widetilde{W}_1$ and $\beta = \widetilde{W}_1^+$. The corresponding current matrix $\widetilde{J}$ takes the form

$$\widetilde{J} = \begin{pmatrix} 0 & \widetilde{W}_2 & \widetilde{W}_{3/2}^{+-} & \widetilde{W}_{3/2}^{++} \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \quad (5.9)$$

The obtained gauged $\mathcal{W}$ -algebra is in this case the algebra $\mathcal{W}[S\ell(4)]$.

Finally, we could have done the reduction from $\mathcal{W}[S\ell(4), S\ell(2)]$ to $\mathcal{W}[S\ell(4)]$ directly by taking the constraints $W_1^- = 1$, $W_{3/2}^{-+} = 1$ and $W_{3/2}^{--} = 0$ which generate the gauge transformation

$$W(x) \longrightarrow \widetilde{W}(x) = W(x) + \int dz \{(\alpha W_{3/2}^{-+} + \beta W_{3/2}^{--} + \gamma W_1^-)(z), W(x)\} + \dots \quad (5.10)$$

The transformed fields are now

$$\begin{aligned} \widetilde{W}_1 &= W_1 + \alpha/2 \\ \widetilde{W}_1^0 &= W_1^0 - \alpha/2 + \gamma \\ \widetilde{W}_1^+ &= W_1^+ - \beta - 2\gamma W_1^0 + \partial\gamma - \gamma^2 + \alpha\gamma/2 \\ \widetilde{W}_2 &= W_2 + 2\partial\alpha + \alpha W_1^0 - 3\alpha W_1 + \beta + \alpha\gamma/2 - \alpha^2 \end{aligned} \quad (5.11)$$

(the expressions for $\widetilde{W}_{3/2}^{+-}$ and $\widetilde{W}_{3/2}^{++}$ are rather heavy). We choose as gauge fixing the conditions $\widetilde{W}_1 = \widetilde{W}_1^0 = \widetilde{W}_1^+ = 0$ which lead to $\alpha = -2W_1$, $\beta = W_1^+ + W_1 \cdot W_1^0 + W_1^0 \cdot W_1^0 - \partial W_1 - \partial W_1^0$, $\gamma = -W_1 - W_1^0$, and the gauge fixed fields $\widetilde{W}_2$, $\widetilde{W}_{3/2}^{+-}$ and $\widetilde{W}_{3/2}^{++}$ generate the algebra $\mathcal{W}[S\ell(4)]$ with the current matrix (5.9).

The other cases can be treated similarly. We summarize in the table 1 the complete secondary reduction scheme in the case of the $\mathcal{W}$ -algebras associated to $S\ell(4)$.

The table 1 is actually completely general. The algebra $\mathcal{W}[S\ell(n), S\ell(2)]$ corresponds always to the less constrained model (i.e. those which contains the most number of fields), while the Abelian $\mathcal{W}[S\ell(n)]$ algebra corresponds to the most constrained model (which contains exactly $n - 1$ fields). From the $\mathcal{W}[S\ell(n), S\ell(2)]$ algebra, one can obtain all the other algebras.

6 Conclusion

We have shown that W -currents can be constrained to provide a new $\mathcal{W}$ -algebra in the same way that conditions on Kac-Moody currents lead to a $\mathcal{W}$ -algebra.

Two $\mathcal{W}$ -algebras can be connected by this mechanism as soon as the set of first and second class constraints of the first algebra is included in the set of constraints of the second one. For $\mathcal{W}[S\ell(n), \mathcal{H}]$ algebras the existence of a suitable gauge, denoted Generalized Horizontal Gauge, insures that $\mathcal{W}(\mathcal{G}, \mathcal{H})$ can be “gauged” to $\mathcal{W}(\mathcal{G}, \mathcal{H}')$ as soon as $\mathcal{H} \subset \mathcal{H}'$. Chains of $\mathcal{W}$ -algebras can then be constructed in relation with the chains of regular, semi-simple




Table 1: Secondary reductions for $\mathcal{W}[Sl(4), \mathcal{H}]$ algebras

subalgebras in $Sl(n)$. Although different, there exists similarity between these chains of subalgebras embeddings and the classification of orbits in the adjoint representation of $\mathcal{G}$: could remarkable geometrical properties known for the $\mathcal{G}$ -orbits [15] be of some relevance for our problem? We have not yet been able to give a pertinent answer to this question.

Let us stress that the above study has been done only for classical $\mathcal{W}$ -algebras. The quantum case needs to be considered. It would also be of some interest to generalize the above results to all series of simple algebras and superalgebras.

Finally, it looks to us specially worthwhile to develop the $\mathcal{W}$ -gauge aspects of this approach. In this respect, the Lagrangian formalism for such a $\mathcal{W}$ -gauge theory is under study and will soon be presented.

References

- [1] L. Fehér, L. O’Raifeartaigh, P. Ruelle, I. Tsutsui and A. Wipf, Phys. Report **222** (1992) *n*° 1, and references therein.
- [2] F.A. Bais, T. Tjin and P. van Driel, Nucl. Phys. **B357** (1991) 632.
- [3] L. Frappat, E. Ragoucy and P. Sorba, Comm. Math. Phys. **157** (1993) 499.
- [4] E. Ragoucy, Nucl. Phys. **B409** (1993) 213.
- [5] A.N. Leznov and M.V. Saveliev, Comm. Math. Phys. **89** (1983) 59;
D.A. Leites, M.V. Saveliev and V.V. Serganova, in *Proceedings of the Third Yurmale Seminar* (VUN Science, Utrecht, The Netherlands, 1986).
- [6] For a general review on W algebras, see P. Bouwknegt and K. Schoutens, Phys. Reports **223** (1993) 183, and references therein.
- [7] F. Delduc, E. Ragoucy and P. Sorba, Comm. Math. Phys. **146** (1992) 403.
- [8] J. de Boer and T. Tjin, Comm. Math. Phys. **158** (1993) 485; id. Comm. Math. Phys. **160** (1994) 317.
- [9] A. Sevrin and W. Troost, Phys. Lett. **B315**, (1993) 304.
A. Sevrin, K. Thielemans and W. Troost, Nucl. Phys. **B407**, (1993) 459.
- [10] J.O. Madsen and E. Ragoucy, *Quantum Hamiltonian Reduction in Superspace Formalism*, Preprint ENSLAPP-A-459/94
- [11] L. Frappat, E. Ragoucy and P. Sorba, Nucl. Phys. **B404** (1993) 805.
- [12] P.A.M. Dirac, *Lecture on Quantum Mechanics*, Academic Press Inc. (NY).
- [13] K. Sundermeyer, *Constrained Dynamics*, Springer-Verlag eds., p.80-81.

- [14] *General properties of classical W algebras*, F. Delduc, L. Frappat, E. Ragoucy and P. Sorba; preprint ENSLAPP-449/93, to appear in Proceedings of *XXII Intern. Conf. on Diff. Geom. Methods in Theor. Physics* (Ixtapa, September 1993)
- [15] L. Michel, Comptes-Rendus Acad. Sci. Paris **272 A** (1971) 433; id. *Statistical Mechanics and Field Theory*, Haifa (Israel) 1971, R.N. Sen and C. Weil eds. (John Wiley and sons, 1972).